

## Supplementary Materials

### Continent-wide drainage reorganization in North America driven by mantle flow

Huilin Wang<sup>1,2,3\*</sup>, Michael Gurnis<sup>1</sup> & Jakob Skogseid<sup>4</sup>

<sup>1</sup> *Seismological Laboratory, California Institute of Technology, Pasadena, CA, USA*

<sup>2</sup> *MOE Key Laboratory of Fundamental Physical Quantities Measurement & Hubei Key*

*Laboratory of Gravitation and Quantum Physics, PGMF and School of Physics, Huazhong*

*University of Science and Technology, Wuhan, 430074, P.R.China*


<sup>3</sup> *Institute of Geophysics and PGM, Huazhong University of Science and Technology, Wuhan,*

*430074, P.R.China*

<sup>4</sup> *Equinor ASA, Fornebu, Norway*

This file contains four supplementary figures

Figures S1, S2, S3 and S4



17


18 **Figure S1. Removed Laramide relief.** All elevation higher than 2 km in the Western

19 Cordillera is removed in the initial model, and added back to surface process models


20 uniformly between 80 to 50 Ma to simulate the Laramide uplift.

21

22



**Figure S2. Paleo-elevation (left panel) and paleo-drainage (right panel) of Models 3 in a North American reference frame.** This model assumes no dynamic topography influences the landscape. In Comparing Model 3 with 1 and 2 we can see that the way the Laramide uplift modified the river patterns is restricted to the western continent, but is not adequate to influence the broader drainage changes.



30


31 **Figure S3. Paleo-elevation maps of Models 4-7 in a North American reference**

32 **frame.** In Model 4, the elastic thickness of continental lithosphere is constant at 40 km;

33 in Model 5, sea level is constant through time; in Model 6, precipitation rate is constant

34 at 1 cm/yr through time; in Model 7, Surface erodibility is fixed at  $1 \times 10^{-7} \text{ yr}^{-1}$ .

35



36

37 **Figure S4. Paleo-drainage of Models 4-7 in a North American reference frame.**

38

39